

Bosonic Structure of a 2-Dimensional Fermion Model with Interaction among Different Species II – N -species Case –

Jiro SAKAMOTO^{*)} and Yasuaki FUKUOKA

Physics department, Shimane University, Matsue 690-8504, Japan

We study a massive Thirring-like model in 2-dimensional space-time, which contains fermions with arbitrary number (N) of different species. This model is an extension of that of a previous paper, where we have considered two-species case. By this extension we expect that we can expose more general structures of this kind of model. We obtain the equivalent boson model with N species to our fermion model. We find that the coupling constant must be set in some regions in order for the model to be physically sensible. It seems hard to find such regions from direct observation of the original fermion model. We also find that for specific values of the coupling constant some of the boson fields disappear from the system. Therefore, the N -species fermion model is described by the boson model with fewer species.

§1. Introduction

The bosonization technique is one of the powerful approaches to study a 2-dimensional fermion system.¹⁾ As Coleman has done in his pioneering work,²⁾ with this technique one can expose hidden properties of such a fermion system though it may be applicable only for charge-zero sectors of fermions, i.e. for sectors of pairs of fermion and anti-fermion. In a previous paper³⁾ (I) using the path-integral method we have derived an equivalent boson model to a two-species fermion model with interaction only among different species. We have found that in order for the model to be physically sensible the coupling constant must be set in some regions. We also found that for a specific value of the coupling constant $g = 2\pi/3$ one boson is decoupled from the system and that the two-species fermion system can be described by the one-species boson model, i.e. the ordinary sine-Gordon model.

In this article we consider the bosonization of the N -species fermion model, which is a direct extension of the model in I. It is an easier but non-trivial extension of the freedom of the bosonization technique. By this extension we expect that we can expose more general structures of this model. As is shown below, we derive the N -species equivalent boson model and find that the coupling constant g must be in the regions $g < -\pi/(N-1)$, $-2\pi/3(N-1) \leq g \leq 2\pi/3$ or $g > \pi$ in order for this boson model to be physically sensible. The region corresponding to the second one was missed in I. We also find that for specific values of the coupling constant some bosons are decoupled from the system, i.e. one or $N-1$ bosons are decoupled for $g = -2\pi/3(N-1)$ or $g = 2\pi/3$, respectively. Therefore, the N -species fermion model is equivalent to $(N-1)$ -species boson model for the former case and to one-species boson model for the latter case.

^{*)} E-mail address: jsakamot@riko.shimane-u.ac.jp

We use the same notations as in I, i.e. in Minkowski space-time, $g_{\mu\nu} = (-1, +1)$ and $\epsilon^{01} = -\epsilon_{01} = 1$. Gamma-matrices are given as $\gamma^0 = -\gamma_0 = i\sigma_x$, $\gamma^1 = \gamma_1 = \sigma_y$, $\gamma_5 = \gamma_0\gamma_1 = \sigma_z$, where σ_x, σ_y and σ_z are the Pauli matrices.

§2. Model

Our initial Lagrangian is given by

$$\mathcal{L} = \sum_{i=1}^N \bar{\psi}_i (\not{\partial} - m) \psi_i + \sum_{i>j=1}^N \frac{1}{2} g j_{i\mu} j_j^\mu, \quad (2.1)$$

where i, j denote the fermion species and

$$j_{i\mu} = i\bar{\psi}_i \gamma_\mu \psi_i. \quad (2.2)$$

In (2.1) a fermion interacts with those of the different species and never with itself directly. When $N = 2$, we can change the sign of g by taking charge conjugation of one of the fermion species. In fact we saw in I that the consequences for $N = 2$ are symmetric for $g \leftrightarrow -g$. On the contrary, for the case $N \geq 3$, there is not such a symmetry, and therefore the sign of g has a physical meaning for this case.

The quartic interaction part in (2.1) is rewritten as

$$\sum_{i>j=1}^N \frac{1}{2} g j_{i\mu} j_j^\mu = \frac{g}{4} \left(\sum_i j_{i\mu} \right)^2 - \frac{g}{4} \sum_i j_{i\mu} j_i^\mu, \quad (2.3)$$

and this is equivalent to

$$\frac{g}{2} \sum_i j_{i\mu} X^\mu - \frac{g}{4} X_\mu X^\mu + \frac{g}{2} \sum_i \left(\frac{1}{2} A_{i\mu} - j_{i\mu} \right) A_i^\mu, \quad (2.4)$$

with using auxiliary vector fields X_μ and $A_{i\mu}$. Now we put

$$B_{i\mu} = X_\mu - A_{i\mu}, \quad (2.5)$$

and integrate out over X_μ by the path-integral formulation to obtain

$$\frac{g}{2} \sum_i j_{i\mu} B_i^\mu - \frac{g}{4(N-1)} \left(\sum_i B_{i\mu} \right)^2 + \frac{g}{4} \sum_i (B_{i\mu})^2. \quad (2.6)$$

Here we should note that for $N \geq 3$ the above expression contains direct coupling among the same species of boson fields $B_{i\mu}$ while fermion fields do not interact directly with themselves in the original Lagrangian (2.1).

In 2-dimensional space-time, we can write vector fields $B_{i\mu}$ with two scalar fields ϕ_i and χ_i as

$$B_{i\mu} = \epsilon_{\mu\nu} \partial^\nu \phi_i + \partial_\mu \chi_i. \quad (2.7)$$

With these scalar fields we transform the fermion fields as

$$\psi_i \rightarrow \psi'_i = \exp \left\{ \frac{ig}{2} (-\gamma_5 \phi_i + \chi_i) \right\} \psi_i, \quad (2.8)$$

to rewrite the Lagrangian as

$$\begin{aligned} \mathcal{L}_{\text{eff}} = & \sum_i \bar{\psi}'_i (\not{\partial} - m \exp\{ig\gamma_5\phi_i\}) \psi'_i - \frac{g}{4(N-1)} \left\{ \left(\sum_i \partial_\mu \chi_i \right)^2 - \left(\sum_i \partial_\mu \phi_i \right)^2 \right\} \\ & + \frac{g}{4} \sum_i (\partial_\mu \chi_i)^2 + \left(\frac{g^2}{4\pi} - \frac{g}{4} \right) \sum_i (\partial_\mu \phi_i)^2. \end{aligned} \quad (2.9)$$

In the above expression the term $(g^2/4\pi)(\partial\phi)^2$ comes from $\det|\exp(-ig\gamma_5\phi_i)|$ in the path-integral measure following Fujikawa.⁴⁾ It is seen that χ_i is decoupled from the other fields and can be integrated out. Then we obtain

$$\begin{aligned} \mathcal{L}_{\text{eff}} = & \sum_i \bar{\psi}_i (\not{\partial} - m \exp\{ig\gamma_5\phi_i\}) \psi_i \\ & + \frac{g}{4(N-1)} \left(\sum_i \partial_\mu \phi_i \right)^2 + \frac{g}{4} \left(\frac{g}{\pi} - 1 \right) \sum_i (\partial_\mu \phi_i)^2, \end{aligned} \quad (2.10)$$

where we write ψ_i for ψ'_i .

The generating functional of the Green functions is given by

$$Z = \int \prod_i d\bar{\psi}_i d\psi_i d\phi_i \exp \left\{ i \int d^2x \mathcal{L}_{\text{eff}} \right\}. \quad (2.11)$$

By the ‘Wick’ rotation we transform ourselves into Euclidean space-time from the Minkowski space-time. Then the generating functional (2.11) is rewritten in Euclidean space-time as

$$Z_E = \int \prod_i d\bar{\psi}_i d\psi_i d\phi_i \exp \left\{ - \int d^2x \mathcal{L}_{\text{eff}} \right\}. \quad (2.12)$$

§3. Perturbative expansion

In order to calculate (2.12) by perturbation theory we choose $\mathcal{L}_0 = \mathcal{L}_{\text{eff}}(m=0)$ of (2.10) as the free Lagrangian. We find that the free fermion and boson propagators are given as

$$\langle \bar{\psi}_i(x) \psi_j(y) \rangle = \frac{\delta_{i,j}}{2\pi} \frac{\gamma \cdot (x-y)}{(x-y)^2}, \quad (3.1)$$

$$\langle \phi_i(x) \phi_j(y) \rangle = \frac{1}{2g} \left\{ \frac{\pi}{(g-\pi)\{\pi+g(N-1)\}} - \frac{\delta_{i,j}}{g-\pi} \right\} \ln(x-y)^2 \mu^2, \quad (3.2)$$

where $\langle \dots \rangle$ denotes the vacuum expectation value of the time-ordered product. Parameter μ is a small infrared cutoff mass, which will be set to zero after the calculations.

Now, we calculate Z_E of (2.12) through the perturbative expansion with respect to m . We note that all the odd order terms vanish because of traceless property of

γ -matrices and the super selection rule for the boson field ϕ , i.e. $\langle \exp i \sum \beta_i \phi \rangle = 0$ unless $\sum \beta_i = 0$ where $\beta_i = \pm g$.^{1), 2)} The $2n$ -th order term of the expansion of Z_E is given as

$$\begin{aligned} Z_E^{(2n)} &= \frac{m^{2n}}{(2n)!} \int \prod d\bar{\psi}_i d\psi_i d\phi_i \left\{ \int dx \sum_{i=1}^N \bar{\psi}_i e^{ig\gamma_5 \phi_i} \psi_i \right\}^{2n} \exp \left\{ - \int dx \mathcal{L}_0 \right\} \\ &= \frac{m^{2n}}{(2n)!} \left\langle \left\{ \int dx \sum_{i=1}^N \bar{\psi}_i e^{ig\gamma_5 \phi_i} \psi_i \right\}^{2n} \right\rangle. \end{aligned} \quad (3.3)$$

Using the identity

$$\bar{\psi} e^{ig\gamma_5 \phi} \psi = e^{ig\phi} \bar{\psi} \Gamma_+ \psi + e^{-ig\phi} \bar{\psi} \Gamma_- \psi, \quad (3.4)$$

where we put $\Gamma_{\pm} = (1 \pm \gamma_5)/2$, we expand (3.3) as

$$\begin{aligned} Z_E^{(2n)} &= m^{2n} \sum_{r_1, r_2, \dots, r_N}^n \frac{\delta_{n, r_1 + r_2 + \dots + r_N}}{(2r_1)!(2r_2)! \dots (2r_N)!} \\ &\quad \times \left\langle \left\{ \int dx \left(e^{ig\phi_1} \bar{\psi}_1 \Gamma_+ \psi_1 + e^{-ig\phi_1} \bar{\psi}_1 \Gamma_- \psi_1 \right) \right\}^{2r_1} \right. \\ &\quad \times \left\{ \int dx \left(e^{ig\phi_2} \bar{\psi}_2 \Gamma_+ \psi_2 + e^{-ig\phi_2} \bar{\psi}_2 \Gamma_- \psi_2 \right) \right\}^{2r_2} \\ &\quad \times \dots \dots \dots \\ &\quad \times \left. \left\{ \int dx \left(e^{ig\phi_N} \bar{\psi}_N \Gamma_+ \psi_N + e^{-ig\phi_N} \bar{\psi}_N \Gamma_- \psi_N \right) \right\}^{2r_N} \right\rangle \\ &= m^{2n} \sum_{r_1, r_2, \dots, r_N}^n \frac{\delta_{n, r_1 + r_2 + \dots + r_N}}{(r_1!)^2 (r_2!)^2 \dots (r_N!)^2} \int \prod dx dy \\ &\quad \times \left\langle \prod_{i=1}^{r_1} \bar{\psi}_1(x_{1,i}) \Gamma_+ \psi_1(x_{1,i}) \bar{\psi}_1(y_{1,i}) \Gamma_- \psi_1(y_{1,i}) \right\rangle \\ &\quad \times \left\langle \prod_{i=1}^{r_2} \bar{\psi}_2(x_{2,i}) \Gamma_+ \psi_2(x_{2,i}) \bar{\psi}_2(y_{2,i}) \Gamma_- \psi_2(y_{2,i}) \right\rangle \\ &\quad \dots \dots \dots \\ &\quad \times \left\langle \exp ig \left[\sum_{i=1}^{r_1} \{ \phi_1(x_{1,i}) - \phi_1(y_{1,i}) \} + \sum_{i=1}^{r_2} \{ \phi_2(x_{2,i}) - \phi_2(y_{2,i}) \} \right. \right. \\ &\quad \left. \left. + \dots \dots \dots + \sum_{i=1}^{r_N} \{ \phi_N(x_{N,i}) - \phi_N(y_{N,i}) \} \right] \right\rangle. \end{aligned} \quad (3.5)$$

With the fermion propagator (3.1) we can calculate a fermion part in the above (3.5)

as

$$\left\langle \prod_{i=1}^{r_k} \bar{\psi}_k(x_{k,i}) \Gamma_+ \psi_k(x_{k,i}) \bar{\psi}_k(y_{k,i}) \Gamma_- \psi_k(y_{k,i}) \right\rangle = \frac{1}{(2\pi)^{2r_k}} \frac{\prod_{i>j}^{r_k} (x_{k,i} - x_{k,j})^2 (y_{k,i} - y_{k,j})^2}{\prod_{i,j}^{r_k} (x_{k,i} - y_{k,j})^2}, \quad (3.6)$$

and we obtain

$$\begin{aligned} Z_E^{(2n)} &= \left(\frac{m}{2\pi} \right)^{2n} \sum_{r_1, r_2, \dots, r_N}^n \frac{\delta_{n, r_1 + r_2 + \dots + r_N}}{(r_1!)^2 (r_2!)^2 \dots (r_N!)^2} \int \prod dx dy \\ &\times \frac{\prod_{i>j}^{r_1} (x_{1,i} - x_{1,j})^2 (y_{1,i} - y_{1,j})^2}{\prod_{i,j}^{r_1} (x_{1,i} - y_{1,j})^2} \times \frac{\prod_{i>j}^{r_2} (x_{2,i} - x_{2,j})^2 (y_{2,i} - y_{2,j})^2}{\prod_{i,j}^{r_2} (x_{2,i} - y_{2,j})^2} \\ &\times \dots \times \frac{\prod_{i>j}^{r_N} (x_{N,i} - x_{N,j})^2 (y_{N,i} - y_{N,j})^2}{\prod_{i,j}^{r_N} (x_{N,i} - y_{N,j})^2} \\ &\times \left\langle \exp ig \left[\sum_{i=1}^{r_1} \{ \phi_1(x_{1,i}) - \phi_1(y_{1,i}) \} + \sum_{i=1}^{r_2} \{ \phi_2(x_{2,i}) - \phi_2(y_{2,i}) \} + \dots \right] \right\rangle. \quad (3.7) \end{aligned}$$

Now, let us calculate the contractions among the boson fields with the same species in the above expression to obtain

$$\begin{aligned} &\left\langle \exp ig \left[\sum_{i=1}^{r_k} \{ \phi_k(x_{k,i}) - \phi_k(y_{k,i}) \} \right] \right\rangle = \exp \left\{ -\frac{g^2 r_k}{2} < \phi_k(0)^2 > \right\} \\ &\times \exp -g^2 \left[\sum_{i>j}^{r_k} \{ < \phi_k(x_{k,i}) \phi_k(x_{k,j}) > + < \phi_k(y_{k,i}) \phi_k(y_{k,j}) > \} - \sum_{i,j}^{r_k} < \phi_k(x_{k,i}) \phi_k(y_{k,j}) > \right] \\ &= \exp \left\{ -\frac{g^2 r_k}{2} < \phi_k(0)^2 > \right\} \\ &\times \exp \left\{ \frac{g^2 (N-1)}{2(g-\pi)\{\pi + g(N-1)\}} \ln \left(\frac{\prod_{i>j}^{r_k} (x_{k,i} - x_{k,j})^2 \mu^2 (y_{k,i} - y_{k,j})^2 \mu^2}{\prod_{i,j}^{r_k} (x_{k,i} - y_{k,j})^2 \mu^2} \right) \right\}, \quad (3.8) \end{aligned}$$

where we have used (3.2). Here we suppose that for a small distance the boson propagator is properly regularized. We find that each fermion part in (3.7) has the same combination of the coordinates $\{x_{k,i}, y_{k,i}\}$ as that in the above expression.

We, therefore, suppose that in the corresponding boson model the free boson propagator gives such a fermion contribution besides the original boson one (3.2) as

$$\begin{aligned} < \phi'_i(x) \phi'_j(y) > = < \phi_i(x) \phi_j(y) > - \frac{\delta_{i,j}}{g^2} \ln(x-y)^2 \mu^2 \\ &= \frac{1}{2g} \left\{ \frac{\pi}{(g-\pi)\{\pi + g(N-1)\}} - \frac{3g-2\pi}{g(g-\pi)} \delta_{i,j} \right\} \ln(x-y)^2 \mu^2, \quad (3.9) \end{aligned}$$

and that potential term is given by

$$m'^2 \{ \cos g \phi'_1 + \cos g \phi'_2 + \dots + \cos g \phi'_N \}. \quad (3.10)$$

The $2n$ -th order term of the perturbative expansion of the generating functional for such a model is calculated as

$$\begin{aligned}
Z_B^{(2n)} &= \frac{m'^{4n}}{(2n)!} \left\langle \left\{ \int dx (\cos g\phi'_1 + \cos g\phi'_2 + \cdots + \cos g\phi'_N) \right\}^{2n} \right\rangle \\
&= m'^{4n} \sum_{r_1, r_2, \dots, r_N}^n \frac{\delta_{n, r_1+r_2+\dots+r_N}}{(2r_1)!(2r_2)! \cdots (2r_N)!} \\
&\quad \times \left\langle \left(\int dx \cos g\phi'_1 \right)^{2r_1} \left(\int dx \cos g\phi'_2 \right)^{2r_2} \cdots \left(\int dx \cos g\phi'_N \right)^{2r_N} \right\rangle \\
&= \left(\frac{m'^2}{2} \right)^{2n} \sum_{r_1, r_2, \dots, r_N}^n \frac{\delta_{n, r_1+r_2+\dots+r_N}}{(r_1!)^2 (r_2!)^2 \cdots (r_N!)^2} \int \prod dx dy \\
&\quad \times \left\langle \exp ig \left[\sum_i^{r_1} \{ \phi'_1(x_{1,i}) - \phi'_1(y_{1,i}) \} + \cdots + \sum_i^{r_N} \{ \phi'_N(x_{N,i}) - \phi'_N(y_{N,i}) \} \right] \right\rangle \\
&= \left(\frac{m'^2}{2} \right)^{2n} \\
&\quad \times \sum_{r_1, \dots, r_N}^n \frac{\delta_{n, r_1+\dots+r_N}}{(r_1!)^2 \cdots (r_N!)^2} \int \prod dx dy \frac{\prod_{i>j}^{r_1} (x_{1,i} - x_{1,j})^2 \mu^2 (y_{1,i} - y_{1,j})^2 \mu^2}{\prod_{i,j}^{r_1} (x_{1,i} - y_{1,j})^2 \mu^2} \times \cdots \\
&\quad \times \left\langle \exp ig \left[\sum_{i=1}^{r_1} \{ \phi_1(x_{1,i}) - \phi_1(y_{1,i}) \} + \cdots + \sum_{i=1}^{r_N} \{ \phi_N(x_{N,i}) - \phi_N(y_{N,i}) \} \right] \right\rangle \\
&\quad \times \exp ng^2 \{ < \phi_i(0)^2 > - < \phi'_i(0)^2 > \}. \tag{3.11}
\end{aligned}$$

If we choose the renormalization of mass as

$$m'^2 = \frac{m\mu}{\pi} \exp \frac{g^2}{2} \{ < \phi_i'^2 > - < \phi_i^2 > \}, \tag{3.12}$$

then Eq.(3.11) is equal to (3.7). From the free boson propagator (3.9) and the potential term (3.10) we find that the boson model with the Lagrangian

$$\mathcal{L}_B = \frac{g^2}{4(3g - 2\pi)} \sum_{i,j} \left[\frac{g}{3g(N-1) + 2\pi} + \frac{g - \pi}{\pi} \delta_{i,j} \right] \partial_\mu \phi'_i \partial^\mu \phi'_j + m'^2 \sum_i \cos g\phi'_i \tag{3.13}$$

is equivalent to the fermion model of (2.1).

§4. Discussion

Let us calculate the eigenvalues of the coefficient matrix $\{A_{i,j}\}$ of the kinetic term in the boson Lagrangian (3.13), which is given by

$$A_{i,j} = \frac{g^2}{2(3g - 2\pi)} \left[\frac{g}{3g(N-1) + 2\pi} + \frac{g - \pi}{\pi} \delta_{i,j} \right]. \tag{4.1}$$

From the equation

$$\det | A_{i,j} - \alpha \delta_{i,j} | = 0 \quad (4.2)$$

we obtain

$$\alpha = \frac{g^2 \{(N-1)g + \pi\}}{2\pi \{3g(N-1) + 2\pi\}} \quad : \text{single root}, \quad (4.3)$$

$$\alpha = \frac{g^2(g - \pi)}{2\pi(3g - 2\pi)} \quad : (N-1)\text{-degenerated roots}. \quad (4.4)$$

In order for the boson model (3.13) to be physically sensible, these eigenvalues must be non-negative, so we have

$$g < -\frac{\pi}{N-1}, \quad (4.5)$$

or

$$-\frac{2\pi}{3(N-1)} \leq g \leq \frac{2\pi}{3}, \quad (4.6)$$

or

$$g > \pi. \quad (4.7)$$

It is hard to find such regions (4.5), (4.6) or (4.7) from direct observation of the fermion model (2.1). In the previous paper I we missed the region corresponding to (4.6). It seems that even in the outside of the regions (4.5), (4.6) and (4.7), we can change the sign of the kinetic part in the Lagrangian by transforming the fields into imaginary ones. By such transformation, however, the potential terms become unbounded from below. This means that in order to fix the physical regions of the coupling constant we must check both kinetic and potential terms in the Lagrangian. The kinetic terms in the Lagrangian (3.13) of the previous paper I look like negative in the region (4.6), but by scaling the boson field we can still define a physically sensible boson model in this region; see below.

At the specific values $g = -\frac{2\pi}{3(N-1)}$ or $g = \frac{2\pi}{3}$ the eigenvalues (4.3) or (4.4) diverge. When $g = -\frac{2\pi}{3(N-1)}$ one boson field corresponding to (4.3) disappears from the system and the original N -species fermion model is described by $(N-1)$ -species boson fields, while for $g = \frac{2\pi}{3}$, $(N-1)$ -boson fields corresponding to (4.4) disappear and the original fermion model is equivalent to the boson model with only one species, i.e. the ordinary sine-Gordon model.

For the case of $N = 2$ the Lagrangian (3.13) in the previous section is rewritten as

$$\begin{aligned} \mathcal{L}_{\mathbf{B}} &= \frac{g^2}{4(3g - 2\pi)} \sum_{i,j}^2 \left[\frac{g}{3g + 2\pi} + \frac{g - \pi}{\pi} \delta_{i,j} \right] \partial_\mu \phi'_i \partial^\mu \phi'_j + m'^2 (\cos g\phi'_1 + \cos g\phi'_2) \\ &= \frac{g^2}{4(3g - 2\pi)} \left[\frac{g}{3g + 2\pi} \{(\partial\phi'_1)^2 + (\partial\phi'_2)^2 + 2\partial\phi'_1 \partial\phi'_2\} + \frac{g - \pi}{\pi} \{(\partial\phi'_1)^2 + (\partial\phi'_2)^2\} \right] \\ &\quad + 2m'^2 \cos \frac{g}{2}(\phi'_1 + \phi'_2) \cos \frac{g}{2}(\phi'_1 - \phi'_2). \end{aligned} \quad (4.8)$$

We put

$$\phi_{\pm} = \frac{g}{2} \sqrt{\frac{g \pm \pi}{\pi(3g \pm 2\pi)}} (\phi'_1 \pm \phi'_2) \quad (4.9)$$

to obtain

$$\begin{aligned} \mathcal{L}_B = & \frac{1}{2} [(\partial\phi_+)^2 + (\partial\phi_-)^2] \\ & + 2m'^2 \cos \left(\sqrt{\frac{\pi(3g+2\pi)}{g+\pi}} \phi_+ \right) \cos \left(\sqrt{\frac{\pi(3g-2\pi)}{g-\pi}} \phi_- \right), \end{aligned} \quad (4.10)$$

which corresponds to the boson Lagrangian in I. As mentioned before, it is seen that the Lagrangian above is invariant for $g \rightarrow -g$ with $\phi_+ \leftrightarrow \phi_-$. There does not exist, however, such a symmetry for the Lagrangian (3.13) when $N \geq 3$.

If $\pi > g > 2\pi/3$ or $-2\pi/3 > g > -\pi$, the potential term in (4.10) becomes

$$2m'^2 \cosh \left(\sqrt{\frac{\pi(3g+2\pi)}{-g-\pi}} \phi_+ \right) \cosh \left(\sqrt{\frac{\pi(3g-2\pi)}{-g+\pi}} \phi_- \right), \quad (4.11)$$

which is unbounded from below, and the model is physically nonsensical. Then g must be set in the region (4.5), (4.6) or (4.7) with $N = 2$.

In our path-integral formulation the massive Thirring-like model is described as the system of massless fermions interacting with massless bosons where the interaction part contains mass parameter. While the bosonization technique may be applicable only for the charge-zero sectors, it is an interesting subject to study the charged sectors, i.e. the bound states of some particles in our formulation.

References

- [1] See for example E. Abdalla, M. C. B. Abdalla and K. D. Rothe, *Non-perturbative methods in two-dimensional quantum field theory* (World Scientific, Singapore, 1991), and M. Stone, *Bosonization* (World Scientific, Singapore, 1994). References are therein.
- [2] S. Coleman, Phys. Rev. **D11** (1975), 2088.
- [3] J. Sakamoto and Y. Heike, Prog. Theor. Phys. **100** (1998), 399.
- [4] K. Fujikawa, Phys. Rev. **D21** (1980), 2848.